

Cite as: W. W. Y. Lau *et al.*, *Science*  
10.1126/science.aba9475 (2020).

# Evaluating scenarios toward zero plastic pollution

Winnie W. Y. Lau<sup>1\*†</sup>, Yonathan Shiran<sup>2\*†</sup>, Richard M. Bailey<sup>3\*†</sup>, Ed Cook<sup>4</sup>, Martin R. Stuchtey<sup>2,5</sup>, Julia Koskella<sup>2</sup>, Costas A. Velis<sup>4\*</sup>, Linda Godfrey<sup>6</sup>, Julien Boucher<sup>7,8</sup>, Margaret B. Murphy<sup>1</sup>, Richard C. Thompson<sup>9</sup>, Emilia Jankowska<sup>2</sup>, Arturo Castillo Castillo<sup>10</sup>, Toby D. Pilditch<sup>3</sup>, Ben Dixon<sup>2</sup>, Laura Koerselman<sup>2</sup>, Edward Kosior<sup>11</sup>, Enzo Favoino<sup>12</sup>, Jutta Gutberlet<sup>13</sup>, Sarah Baulch<sup>1</sup>, Meera E. Atreya<sup>2</sup>, David Fischer<sup>2</sup>, Kevin K. He<sup>1</sup>, Milan M. Petit<sup>2</sup>, U. Rashid Sumaila<sup>14</sup>, Emily Neil<sup>3</sup>, Mark V. Bernhofen<sup>4</sup>, Keith Lawrence<sup>1</sup>, James E. Palardy<sup>1\*†</sup>

<sup>1</sup>The Pew Charitable Trusts, 901 E Street NW, Washington, DC 20004, USA. <sup>2</sup>SYSTEMIQ, 69 Carter Lane, London EC4V 5EQ, UK. <sup>3</sup>School of Geography and the Environment, Oxford University, Oxford OX1 3QY, UK. <sup>4</sup>School of Civil Engineering, University of Leeds, Leeds LS2 9JT, UK. <sup>5</sup>Institute of Geography, University of Innsbruck, Innrain 52, 6020 Innsbruck, Austria. <sup>6</sup>Council for Scientific and Industrial Research, Pretoria 0001, South Africa. <sup>7</sup>EA—Shaping Environmental Action, Ch. des vignes d'argent 7, CH 1004 Lausanne, Switzerland. <sup>8</sup>University of Applied Sciences and Arts Western Switzerland//HES-SO, HEIG-VD, Yverdon-les-Bains, Switzerland. <sup>9</sup>School of Biological and Marine Sciences, University of Plymouth, Plymouth, PL4 8AA, UK. <sup>10</sup>Faculty of Natural Sciences, Centre for Environmental Policy, Imperial College London, London SW7 2AX, UK. <sup>11</sup>Nextek, 1 Kensington Gore London SW7 2AR, UK. <sup>12</sup>Scuola Agraria del Parco di Monza, Viale Cavriga 3 20900 Monza (MB), Italy. <sup>13</sup>Department of Geography, University of Victoria, Victoria, British Columbia, V8W 2Y2, Canada. <sup>14</sup>Institute for the Oceans and Fisheries and School of Public Policy and Global Affairs, University of British Columbia, Vancouver, British Columbia, V6T 1Z4, Canada.

\*Corresponding author. Email: [wlau@pewtrusts.org](mailto:wlau@pewtrusts.org) (W.W.Y.L.); [yonathan.shiran@systemiq.earth](mailto:yonathan.shiran@systemiq.earth) (Y.S.); [richard.bailey@ouce.ox.ac.uk](mailto:richard.bailey@ouce.ox.ac.uk) (R.M.B.); [c.velis@leeds.ac.uk](mailto:c.velis@leeds.ac.uk) (C.A.V.); [jpalardy@pewtrusts.org](mailto:jpalardy@pewtrusts.org) (J.E.P.)

†These authors contributed equally to this work.

**Plastic pollution is a pervasive and growing problem. To estimate the effectiveness of interventions to reduce plastic pollution, we modeled stocks and flows of municipal solid waste and four sources of microplastics through the global plastic system for five scenarios between 2016 and 2040. Implementing all feasible interventions reduced plastic pollution by 40% from 2016 rates and 78% relative to 'business as usual' in 2040. Even with immediate and concerted action, 710 million metric tons of plastic waste cumulatively entered aquatic and terrestrial ecosystems. To avoid a massive build-up of plastic in the environment, coordinated global action is urgently needed to reduce plastic consumption, increase rates of reuse, waste collection and recycling, expand safe disposal systems and accelerate innovation in the plastic value chain.**

Plastic pollution is globally ubiquitous. It is found throughout the oceans, in lakes and rivers, in soils and sediments, in the atmosphere, and in animal biomass. This proliferation has been driven by rapid growth in plastic production and use combined with linear economic models that ignore the externalities of waste (1, 2). A sharp rise in single-use plastic consumption and an expanding 'throw-away' culture (1) have exacerbated the problem. Waste management systems do not have sufficient capacity at the global level to safely dispose of or recycle waste plastic (3, 4), resulting in an inevitable increase in plastic pollution into the environment. Previous studies estimated approximately 8 million metric tons (Mt) of macroplastic (5) and 1.5 Mt of primary microplastic (6) enter the ocean annually. Comparable estimates for terrestrial plastic pollution have yet to be quantified. If plastic production and waste generation continue to grow at current rates, the annual mass of mismanaged waste has been projected to more than double by 2050 (1, 2) and the cumulative mass of ocean plastic could increase by an order of magnitude from 2010 levels by 2025 (5). Despite the magnitude of these flows, the efficacy and economic costs of solutions proposed to solve the plastic waste problem – the uncontrolled release of plastic

waste into the environment resulting from ineffective management – remains unknown.

A growing body of evidence points to a broad range of detrimental effects of plastic pollution. Nearly 700 marine species and over 50 freshwater species are known to have ingested or become entangled in macroplastic (7, 8) and there is growing evidence that plastic is ingested by a wide range of terrestrial organisms (9). Plastic pollution impacts many aspects of human well-being: affecting the aesthetics of beaches (10), blocking drainage and wastewater engineering systems (11) and providing a breeding ground for disease vectors (10, 12). The lower-bound estimate of the economic costs of plastic pollution on fishing, tourism and shipping have been estimated at USD 13 billion annually (13). Although harmful effects of microplastic (here defined as plastics < 5 mm) have not been consistently demonstrated, ingestion has been documented across trophic levels and at all depths of the ocean in individual organisms and species assemblages (8, 14) and in terrestrial organisms (15). Microplastics are also increasingly found in the human food system though their impacts on human health are difficult to assert and require further research (16, 17). Plastic production, collection and

disposal are also major sources of greenhouse gas (GHG) emissions (18).

Cost-effective solutions to managing plastic waste vary considerably across geographies and social settings (3), and a variety of solutions to the plastic pollution problem have been proposed at local, national and regional levels (19, 20). Some proposed interventions focus on post-consumption management, requiring considerable growth in investment and capacity of waste management solutions (21, 22). Other interventions prioritize reducing plastic through replacement with alternative products, reuse, and the development of new delivery models (23). Individual countries have established bans or levies on select plastic products, with a particular focus on banning single-use carrier bags and microbeads in cosmetic products (24, 25). The European Union recently adopted a directive on single-use plastics (26) while the Basel Convention was amended to regulate the international trade of plastic waste (27). The scientific community and non-governmental organizations are also working to identify solutions (21, 28). Despite these efforts, a global evidence-based strategy that includes practical and measurable interventions aimed at reducing plastic pollution does not yet exist.

### Modeling approach

Designing an effective global strategy requires an understanding of the mitigation potential of different solutions and the magnitude of global effort needed to appreciably reduce plastic pollution. To estimate mitigation potential under different intervention scenarios, we developed the Plastics-to-Ocean (P<sub>2</sub>O) model. P<sub>2</sub>O is a data-driven coupled ordinary differential equation (ODE) model that calculates the flow of plastics through representative systems. We used the model to characterize key stocks and flows for land-based sources of plastic pollution across the entire value chain for municipal solid waste (MSW) macroplastics (figs. S1 and S2) and four sources of primary microplastics (i.e., those entering the environment as microplastics) [supplementary materials (SM) section 15 and figs. S3 to S6). Crucially, it provides estimates of plastic waste input into the environment. Costs are calculated as a function of modeled plastic flows, and changes in costs due to production scale and technological advancement are accounted for through learning curves and returns to scale (SM section 16.1).

Projected growth in demand for plastic was calculated using country-level population size (29), per capita macroplastic MSW (30, 31) and microplastic-generating product use and loss rates. Per capita waste generation and waste management processes (e.g., collection costs, collection and processing rates, recycling recovery value) and rates of primary microplastic generation vary by geography and plastic category/source (6, 32–34). To account for these differences, the global population was split across eight geographic

archetypes based on World Bank income categories (low income, lower- and upper- middle income and high income); and United Nations urban-rural classifications (29). Populations were further differentiated by their distance to water (< 1 km or > 1 km) to estimate their relative flows of plastic pollution to terrestrial versus aquatic (lakes, rivers and marine environments) systems. To account for different waste management pathways (35) and movement rates of waste in the environment (35), MSW plastics were differentiated into three material categories: rigid monomaterial, flexible monomaterial and multi-material/multi-layer. Four microplastic sources were modeled: synthetic textiles, tires, plastic pellets and personal care products.

Five scenarios were developed to estimate reductions in plastic pollution over the period 2016–2040. Scenarios were defined by four high-level classes of interventions (reduce, substitute, recycle, dispose) and eight system interventions: (i) reducing plastic quantity in the system, (ii) substituting plastics with alternative materials and delivery systems, (iii) implementing design for recycling, (iv) increasing collection capacity, (v) scaling-up sorting and mechanical recycling capacity, (vi) scaling-up chemical conversion capacity, (vii) reducing post-collection environmental leakage, and (viii) reducing trade in plastic waste (table S7). Scenarios modeled include: (i) ‘Business as Usual’ (BAU), (ii) ‘Collect and Dispose’, (iii) ‘Recycling’, (iv) ‘Reduce and Substitute’, and (v) an integrated ‘System Change’ scenario that implemented the entire suite of interventions (tables S8 and S57).

At all relevant geographical scales, waste production and handling data are notoriously difficult to obtain. Many model inputs have a high degree of uncertainty which was propagated using Monte Carlo sampling. Data inputs and assigned uncertainties are described in supplemental material (SM section 5.6). In the absence of datasets with which to formally validate the model, sensitivity analyses were conducted to quantify the influence of individual model inputs and to identify key drivers of plastic pollution. Model outputs from the BAU scenario were also compared against results from other global studies (2, 5, 36).

### Business as usual

The BAU scenario highlights the scale of the plastic pollution problem and provides a baseline from which to compare alternative intervention strategies (Fig. 1). At a global scale from 2016–2040, the annual rate of macro- and microplastic entering aquatic systems from land increased 2.6-fold (Fig. 1C and Table 1). Over the same period, the rate of plastic pollution retained in terrestrial systems increased 2.8-fold (Fig. 1D and Table 1).

When current commitments to reducing plastic pollution were modeled assuming full implementation (SM section 9.1), annual plastic pollution rates into aquatic and terrestrial

environments decreased by only 6.6% [95% Confidence Interval: 5.4, 7.9] (37) and 7.7% [5.2, 10] by 2040, respectively (Fig. 1A). This result confirms that current commitments coupled with appropriate policies can reduce plastic waste input into the environment but also shows that considerable additional effort will be needed to match the unprecedented scale of projected environmental plastic pollution.

Plastic pollution rates were found to be particularly sensitive to total plastic mass, collection rates, and the ratio of managed to mismanaged waste. For example, a 1 t reduction in plastic MSW mass (i.e., through reduce and substitute interventions) decreased aquatic plastic pollution by an average of 0.088 t in low and middle-income archetypes and an average of 0.0050 t in high-income archetypes. Across all archetypes, an equivalent increase in the collection of plastic waste (through formal and informal sectors) resulted in an average 0.18 t decrease in aquatic plastic pollution, while a similar decrease in post-collection mismanaged waste produced an average 0.10 t decrease in aquatic plastic pollution.

### Scenarios to reduce plastic pollution

The focus of plastic pollution reduction strategies can be broadly partitioned into upstream (pre-consumption, e.g., reducing demand) and downstream (post-consumption, e.g., collection and recycling) measures. To parameterize the development of waste management and recycling solutions in the ‘Collect and Dispose’, ‘Recycling’, and ‘System Change’ scenarios, we estimated maximum foreseen growth and implementation rates based on historical trends and expert panel consensus assessment (SM section 1). In a limited number of cases where data were not available in the published literature, we conducted interviews with industry experts or purchased proprietary data from industry market research databases. Compared to BAU, the annual combined terrestrial and aquatic plastic pollution rates were reduced by 57% in 2040 [45, 69] under the ‘Collect and Dispose’ scenario, and by 45% [35, 54] under the ‘Recycling’ scenario (Fig. 1, A and B).

Strategies focused on upstream (pre-consumption) solutions were represented by the ‘Reduce and Substitute’ scenario. We developed a feasibility assessment framework to model the potential development of upstream solutions aimed at reducing the volume of plastics used and disposed of into the waste stream (SM section 9). Fifteen major plastic applications were assessed against four criteria for technology readiness and unintended consequences related to health/food safety, consumer acceptance (e.g., convenience, climate change impacts) and affordability (tables S21 and S22). The feasibility of substitution with alternative material was assessed against the potential for scaling to meaningful levels within the modeling period. Paper, coated paper and compostable materials met these criteria. Under the ‘Reduce

and Substitute’ scenario, annual combined terrestrial and aquatic plastic pollution in 2040 decreased 59% [47, 72] relative to BAU while annual plastic production decreased by 47% [44, 49]. Consequently, plastic production in 2040 under the ‘Reduce and Substitute’ scenario (220 Mt/y [200, 240]) was similar to production in 2016 (210 Mt/y [200, 230]).

Neither pre- nor post-consumption interventions alone are sufficient to address the plastic problem. Combining the maximum foreseen application of pre- and post-consumption solutions represents the most aggressive possible solution given current technology: the ‘System Change’ scenario. In this scenario, annual combined terrestrial and aquatic plastic pollution decreased by 78% [62, 94] relative to BAU in 2040, but only by 40% [31, 48] relative to 2016 pollution rates (Fig. 1, A and B, and Table 1). In 2040, the annual rate of land-based sources of plastic entering aquatic and terrestrial systems decreased by 82% [68, 95] and 76% [55, 97] relative to BAU, respectively (Fig. 1, C and D, and Table 1).

Under the ‘System Change’ scenario in 2040, a substantial reduction in mismanaged and disposed waste was achieved through increases in the proportion of plastic demand reduced, substituted by alternative materials and recycled (Fig. 2A, and Table 1). These changes to the plastic system resulted in 11% [10, 12] less virgin plastic being produced in 2040 under the ‘System Change’ scenario than was produced in 2016, and 55% [51, 58] less than in 2040 under BAU. Moreover, this reduction was driven by increases in recycled plastic feedstock, which have lower life-cycle GHG emissions (18). Taken together, the ‘System Change’ scenario moves toward achieving a circular economy in which resources are conserved, waste generation is minimized (38) and GHG emissions reduced.

The present value of cumulative, global waste management operations from 2016 to 2040 was approximated to assess the relative cost of each scenario (Fig. 3). Among scenarios, costs varied by less than 20% relative to BAU, were lowest under the ‘System Change’ and ‘Recycling’ scenarios, and highest for the ‘Collect and Dispose’ scenario. Costs under the ‘System Change’ scenario were 18% [14, 23] lower than BAU, with increased waste management costs offset by costs savings from reduced plastic production and revenues from recyclate sales, which increased due to product redesign and improved economics of recycling (SM section 16.8). These costs represent only waste management costs, which are generally borne by taxpayers. Corporate engagement, through improved product design, alternative material development and new business models will be necessary to achieve pollution levels observed in the ‘System Change’ scenario. This engagement will likely require a significant shift in private sector investment.



Our results underline the urgency with which extensive interventions are needed. Despite a considerable reduction in annual plastic production and an increase in the proportion of MSW that is effectively managed under the best-case ‘System Change’ scenario, a substantial amount of plastic waste remained mismanaged (i.e., not collected and sorted, recycled or safely disposed) between 2016 and 2040. When implementation of interventions begins in 2020, the cumulative mass of plastic pollution added between 2016 and 2040 amounts to 250 Mt [190, 310] in aquatic systems (Fig. 4A) and 460 Mt [300, 640] in terrestrial systems (Fig. 4B), approximately 1 and 2 times the total annual plastic production in 2016, respectively. If implementation of interventions is delayed by only 5 years, an additional 300 Mt of mismanaged plastic waste is expected to accumulate in the environment.

### Outlook by plastic category

The complex composition of multi-material plastics limits the technical feasibility of sorting and reprocessing (39), decreasing the economic attractiveness of recycling. Accordingly, the annual production of these plastics decreased by 19 Mt [18, 20] from 2016 to 2040 under the ‘System Change’ scenario, with a shift of similar magnitude to flexible mono-material plastic production (20 Mt/y [19, 21]).

Due to the relative ease of collection and sorting, recycling was dominated by rigid plastics in all archetypes and across all scenarios (Fig. 4C). Under the ‘System Change’ scenario in 2040, rigid plastics represented 62% [58, 67] of the annual mass of recycling, with a sizeable component of flexible mono-material plastic (33% [28, 37]) (Fig. 5A). In comparison, only 5.0% [4.2, 5.4] of recycled material was derived from multi-material/multilayer waste plastic (Fig. 5A).

The diversity of polymer types, surface contamination and low density of post-consumer flexible monomaterial limit their capacity for recycling, particularly in geographies where waste collection services are provided by the informal sector. At a global scale, the absolute and relative contribution of flexible monomaterial plastics to environmental pollution grew between 2016 and 2040, from 45% [35, 56] to 56% [40, 73] in aquatic environments and from 37% [18, 52] to 48% [22, 67] in terrestrial environments (Fig. 5, B and C). Accordingly, finding an economically viable solution to effectively manage flexible plastics will be essential for solving the plastic pollution problem.

Similarly, the proportion of total plastic pollution originating from microplastics in the ‘System Change’ scenario grew from 11% [6.5, 18] to 23% [11, 42] in aquatic systems and from 16% [8.2, 27] to 31% [18, 51] in terrestrial systems over the modeled period (Fig. 5, B and C). Technologies to capture microplastics, which often rely on stormwater and wastewater management and treatment, are rarely economically feasible – even in wealthy regions – due to associated

infrastructure costs. This technological challenge is particularly acute for tire particles, which contributed 93% [83, 96] of global microplastic pollution by mass in 2040.

### Difficulties to overcome

Scaling collection to all households at a global level is a monumental task that would require connecting over a million additional households to MSW collection services per week from 2020 to 2040; the majority of these unconnected households are in middle-income countries. The effort to increase household waste collection will therefore require a key role for ‘waste pickers’ (the informal collection and recycling sector (40)), who link the service chain (MSW collection) to the value chain (recycling) in low- and middle-income settings. Globally, this sector was responsible for 58% [55, 64] of post-consumer plastic waste collected for recycling in 2016. To incentivize the collection of low-value plastics (flexible monomaterial and multimaterial / multilayer plastic) by the informal sector, the profitability of recycling these materials would need to rise to create demand for their collection. Accordingly, investments in collection infrastructure must be coordinated with improved governance around collection, sorting and safe management of generated waste (41).

Mismanaged plastic waste (i.e., in dumpsites, openly burned or released into aquatic or terrestrial environments) is associated with a range of risks to human and ecological health (42). Substantial quantities of such waste are likely to continue to be emitted into the environment or openly burned through time. Under the ‘System Change’ scenario, in addition to aquatic and terrestrial pollution, approximately 250 Mt [130, 380] of waste plastic would accumulate in open dumpsites from 2016 to 2040 and remain a potential source of environmental pollution (Fig. 4D). Many communities in emerging economies with inadequate waste management services and infrastructure burn waste residentially or in open dumpsites without emissions controls. Open burning transfers the pollution burden to air, water and land via the generation of GHGs, particulate matter (including microplastic particles) and harmful chemicals such as dioxins and other persistent organic pollutants (43, 44). Despite its human health and environmental consequences, open burning was the single largest component of mismanaged plastic waste under all scenarios, with 1200 Mt [940, 1400] of plastic burned in the ‘System Change’ scenario between 2016 and 2040 (Fig. 4D). It therefore remains a stubborn pollution and social justice problem in need of an effective solution.

Though not strictly mismanaged, the net export of waste from high-income to upper- and lower-middle income countries grew from 2.7 Mt/y [2.4, 4.7] in 2016 to 3.8 Mt/y [0.7, 7.2] in 2040 under BAU. Though a comparatively small amount, these exports have the potential to increase the fraction of mismanaged plastic waste, as receiving countries often have

insufficient capacity to manage their own waste. Consequently, importing waste for recycling can have the unintended consequence of displacing these developing economies' capacity to recycle their domestic waste (45).

Although efforts to measure the amount of plastic pollution entering rivers and the ocean have increased in recent years (46–48), key data gaps remain. To better estimate the effects of consumer, corporate and policy actions on solving the plastic pollution problem, additional empirical data are needed throughout the plastics system – particularly in developing economies. Moreover, a more complete accounting of the benefits, costs and externalities of plastic use is needed to design policies which align social and financial incentives and minimize plastic pollution. These data deficiencies currently prevent application of the model at finer geographical scales and limit the granularity of the system representation. In particular, data from the informal sector of the global waste management system are scarce, as are data which shed light on the importance of post-collection MSW mismanagement. Additional quantitative data are also needed to better understand key sources, rates and pathways for microplastic pollution and for maritime sources of plastic pollution.

### Addressing the plastic pollution problem

Our analysis indicates that urgent and coordinated action combining pre- and post-consumption solutions could reverse the increasing trend of environmental plastic pollution. While no silver bullet exists, 78% of the plastic pollution problem can be solved by 2040 using current knowledge and technologies and at a lower net cost for waste management systems compared to BAU. However, with long degradation times, even a 78% reduction from BAU pollution rates results in a massive accumulation of plastic waste in the environment. Moreover, even if this system change is achieved, plastic production and unsound waste management activities will continue to emit large quantities of GHGs. Further innovation in resource-efficient and low-emission business models, reuse and refill systems, sustainable substitute materials, waste management technologies and effective government policies are needed. Such innovation could be financed by re-directing existing and future investments in virgin plastic infrastructure. Substantial commitments to improving the global plastic system are required from businesses, governments and the international community to solve the ecological, social and economic problems of plastic pollution and achieve near-zero input of plastics into the environment.

### REFERENCES AND NOTES

1. R. Geyer, J. R. Jambeck, K. L. Law, Production, use, and fate of all plastics ever made. *Sci. Adv.* **3**, e1700782 (2017). [doi:10.1126/sciadv.1700782](https://doi.org/10.1126/sciadv.1700782) [Medline](#)
2. L. Lebreton, A. Andrady, Future scenarios of global plastic waste generation and disposal. *Palgrave Commun.* **5**, 6 (2019). [doi:10.1057/s41599-018-0212-7](https://doi.org/10.1057/s41599-018-0212-7)
3. C. A. Velis, D. Lerpiniere, M. Tsakona, "Prevent marine plastic litter - now!" (International Solid Waste Association, Vienna, 2017).
4. D. C. Wilson, C. A. Velis, Waste management—still a global challenge in the 21st century: An evidence-based call for action. *Waste Manag. Res.* **33**, 1049–1051 (2015). [doi:10.1177/0734242X15616055](https://doi.org/10.1177/0734242X15616055) [Medline](#)
5. J. R. Jambeck, R. Geyer, C. Wilcox, T. R. Siegler, M. Perryman, A. Andrady, R. Narayan, K. L. Law, Plastic waste inputs from land into the ocean. *Science* **347**, 768–771 (2015). [doi:10.1126/science.1260352](https://doi.org/10.1126/science.1260352) [Medline](#)
6. J. Boucher, D. Friot, "Primary microplastics in the oceans: a global evaluation of sources" (IUCN, Gland, Switzerland, 2017).
7. S. C. Gall, R. C. Thompson, The impact of debris on marine life. *Mar. Pollut. Bull.* **92**, 170–179 (2015). [doi:10.1016/j.marpolbul.2014.12.041](https://doi.org/10.1016/j.marpolbul.2014.12.041) [Medline](#)
8. C. M. Rochman, M. A. Browne, A. J. Underwood, J. A. van Franeker, R. C. Thompson, L. A. Amaral-Zettler, The ecological impacts of marine debris: Unraveling the demonstrated evidence from what is perceived. *Ecology* **97**, 302–312 (2016). [doi:10.1890/14-2070.1](https://doi.org/10.1890/14-2070.1) [Medline](#)
9. E. Huerta Lwanga, J. Mendoza Vega, V. Ku Quej, J. L. A. Chi, L. Sanchez Del Cid, C. Chi, G. Escalona Segura, H. Gertsen, T. Salánki, M. van der Ploeg, A. A. Koelmans, V. Geissen, Field evidence for transfer of plastic debris along a terrestrial food chain. *Sci. Rep.* **7**, 14071 (2017). [doi:10.1038/s41598-017-14588-2](https://doi.org/10.1038/s41598-017-14588-2) [Medline](#)
10. K. J. Wyles, S. Pahl, K. Thomas, R. C. Thompson, Factors that can undermine the psychological benefits of coastal environments: Exploring the effect of tidal state, presence, and type of litter. *Environ. Behav.* **48**, 1095–1126 (2016). [doi:10.1177/0013916515592177](https://doi.org/10.1177/0013916515592177) [Medline](#)
11. J. Fobil, J. Hogarth, The dilemmas of plastic wastes in a developing economy: Proposals for a sustainable management approach for Ghana. *West Afr. J. Appl. Ecol.* **10**, 1 (2009). [doi:10.4314/wajae.v10i1.45716](https://doi.org/10.4314/wajae.v10i1.45716)
12. E. Boelee, G. Geerling, B. van der Zaan, A. Blauw, A. D. Vethaak, Water and health: From environmental pressures to integrated responses. *Acta Trop.* **193**, 217–226 (2019). [doi:10.1016/j.actatropica.2019.03.011](https://doi.org/10.1016/j.actatropica.2019.03.011) [Medline](#)
13. United Nations Environment Programme, "Valuing plastics: the business case for measuring, managing and disclosing plastic use in the consumer goods industry" (2014).
14. D. S. Green, B. Boots, D. J. Blockley, C. Rocha, R. Thompson, Impacts of discarded plastic bags on marine assemblages and ecosystem functioning. *Environ. Sci. Technol.* **49**, 5380–5389 (2015). [doi:10.1021/acs.est.5b00277](https://doi.org/10.1021/acs.est.5b00277) [Medline](#)
15. A. A. de Souza Machado, W. Kloas, C. Zarfl, S. Hempel, M. C. Rillig, Microplastics as an emerging threat to terrestrial ecosystems. *Glob. Change Biol.* **24**, 1405–1416 (2018). [doi:10.1111/gcb.14020](https://doi.org/10.1111/gcb.14020) [Medline](#)
16. L. G. A. Barboza, A. Dick Vethaak, B. R. B. O. Lavorante, A. K. Lundebye, L. Guilhermino, Marine microplastic debris: An emerging issue for food security, food safety and human health. *Mar. Pollut. Bull.* **133**, 336–348 (2018). [doi:10.1016/j.marpolbul.2018.05.047](https://doi.org/10.1016/j.marpolbul.2018.05.047) [Medline](#)
17. D. Peixoto, C. Pinheiro, J. Amorim, L. Oliva-Teles, L. Guilhermino, M. N. Vieira, Microplastic pollution in commercial salt for human consumption: A review. *Estuar. Coast. Shelf Sci.* **219**, 161–168 (2019). [doi:10.1016/j.ecss.2019.02.018](https://doi.org/10.1016/j.ecss.2019.02.018)
18. J. Zheng, S. Suh, Strategies to reduce the global carbon footprint of plastics. *Nat. Clim. Chang.* **9**, 374–378 (2019). [doi:10.1038/s41558-019-0459-z](https://doi.org/10.1038/s41558-019-0459-z)
19. D. Xanthos, T. R. Walker, International policies to reduce plastic marine pollution from single-use plastics (plastic bags and microbeads): A review. *Mar. Pollut. Bull.* **118**, 17–26 (2017). [doi:10.1016/j.marpolbul.2017.02.048](https://doi.org/10.1016/j.marpolbul.2017.02.048) [Medline](#)
20. J. K. Abbott, U. R. Sumaila, Reducing marine plastic pollution: Policy insights from economics. *Rev. Environ. Econ. Policy* **13**, 327–336 (2019). [doi:10.1093/reep/rez007](https://doi.org/10.1093/reep/rez007)
21. M. Cordier, T. Uehara, How much innovation is needed to protect the ocean from plastic contamination? *Sci. Total Environ.* **670**, 789–799 (2019). [doi:10.1016/j.scitotenv.2019.03.258](https://doi.org/10.1016/j.scitotenv.2019.03.258) [Medline](#)

22. I. E. Napper, R. C. Thompson, Environmental deterioration of biodegradable, oxo-biodegradable, compostable, and conventional plastic carrier bags in the sea, soil, and open-air over a 3-year period. *Environ. Sci. Technol.* **53**, 4775–4783 (2019). [doi:10.1021/acs.est.8b06984](https://doi.org/10.1021/acs.est.8b06984) [Medline](#)
23. Ellen MacArthur Foundation, “The New Plastics Economy: rethinking the future of plastics” (2016).
24. R. E. J. Schnurr, V. Alboiu, M. Chaudhary, R. A. Corbett, M. E. Quanz, K. Sankar, H. S. Srain, V. Thavarajah, D. Xanthos, T. R. Walker, Reducing marine pollution from single-use plastics (SUPs): A review. *Mar. Pollut. Bull.* **137**, 157–171 (2018). [doi:10.1016/j.marpolbul.2018.10.001](https://doi.org/10.1016/j.marpolbul.2018.10.001) [Medline](#)
25. P. Dauvergne, The power of environmental norms: Marine plastic pollution and the politics of microbeads. *Env. Polit.* **27**, 579–597 (2018). [doi:10.1080/09644016.2018.1449090](https://doi.org/10.1080/09644016.2018.1449090)
26. European Union, “Directive (EU) 2019/904 of the European Parliament and of the Council of 5 June 2019 on the reduction of the impact of certain plastic products on the environment.” (2019).
27. United Nations Environment Programme Secretariat of the Basel Convention, “Decision BC-14/13: further actions to address plastic waste under the Basel Convention” (Châtelaine GE, Switzerland, 2018).
28. F. Oosterhuis, E. Papyrakis, B. Boteler, Economic instruments and marine litter control. *Ocean Coast. Manage.* **102**, 47–54 (2014). [doi:10.1016/j.ocecoaman.2014.08.005](https://doi.org/10.1016/j.ocecoaman.2014.08.005)
29. United Nations Department of Economic and Social Affairs Population Division, “World Urbanization Prospects: The 2018 Revision (ST/ESA/SER.A/420)” (United Nations, New York: United Nations, 2019).
30. S. Kaza, L. Yao, P. Bhada-Tata, F. Van Woerden, *What a Waste 2.0: A Global Snapshot of Solid Waste Management to 2050* (The World Bank, Washington, DC, 2018).
31. M. Economics, “The Circular Economy - A powerful force for climate mitigation” (Stockholm, 2018).
32. OECD, *Environment at a Glance 2015: OECD Indicators* (OECD Publishing, Paris, 2015).
33. WRAP, “Defining what’s recyclable and best in class polymer choices for packaging” (v1, Banbury, 2019).
34. E. G. Ashton, W. Kindlein Jr., R. Demori, L. H. A. Cândido, R. Mauler, Recycling polymeric multi-material products through micronization. *J. Clean. Prod.* **116**, 268–278 (2016). [doi:10.1016/j.jclepro.2016.01.018](https://doi.org/10.1016/j.jclepro.2016.01.018)
35. OECD, “Improving markets for recycled plastics” (OECD Publishing, Paris, France, 2018).
36. Excluding microplastic pollution, the P<sub>2</sub>O model estimates that 9.8 Mt/y [7.7, 12] of plastic pollution enters aquatic systems in 2016 and 16 Mt/y [12, 20] in 2025. These outputs closely align with ranges reported by Jambeck *et al.* (5), who report a midpoint of 9.1 Mt/y (25% of mismanaged waste entering the ocean) [5.5 Mt/y for 15%; 14.6 Mt/y for 40%] in 2015 and 17.5 Mt/y [10.5 Mt/y, 28 Mt/y] in 2025. Estimated masses of mismanaged waste reported here (for 2016: 87 Mt [81, 93]; for 2020: 108 Mt [101, 126]) are higher than, but in the same order of magnitude, as those reported by Jambeck *et al.* (5) (2016: 36.5 Mt; 2020: 69.9 Mt). This is unsurprising, as Jambeck *et al.* (5) do not estimate mismanaged waste generated > 50 km from the coast. In early model time steps, estimated mismanaged MSW presented here (2016: 87 Mt [81, 93]; 2020: 108 Mt [101, 126]) align well with those presented by Lebreton and Andrady (2) (2015: 80 Mt [60, 99]; “2020: 96 Mt [75, 115]). Estimates of mismanaged MSW from the two models diverge into the future: we estimated 228 Mt [213, 252] in 2040 while Lebreton and Andrady (2) estimated 155 Mt [118, 188]. This divergence may be due to several differences among the models in urban/rural population splits and constraints on waste management and recycling capacities applied in P<sub>2</sub>O.
37. Reported results represent model output when inputs are assumed to have no error. 95% Confidence Intervals (CI) were calculated using Monte Carlo sampling (see SM section 5.6). Hereafter, values in square brackets represent the lower and upper bounds of the CI.
38. M. Crippa *et al.*, “A circular economy for plastics – insights from research and innovation to inform policy and funding decisions” (European Commission, Brussels, Belgium, 2019).
39. S. Slater, T. Crichton, “Recycling of laminated packaging” (Banbury, UK, 2011).
40. C. A. Velis, D. C. Wilson, O. Rocca, S. R. Smith, A. Mavropoulos, C. R. Cheeseman, An analytical framework and tool (‘InteRa’) for integrating the informal recycling sector in waste and resource management systems in developing countries. *Waste Manag. Res.* **30** (Suppl), 43–66 (2012). [doi:10.1177/0734242X12454934](https://doi.org/10.1177/0734242X12454934) [Medline](#)
41. D. C. Wilson, L. Rodic, M. J. Cowing, C. A. Velis, A. D. Whiteman, A. Scheinberg, R. Vilches, D. Masterson, J. Stretz, B. Oelz, ‘Wasteaware’ benchmark indicators for integrated sustainable waste management in cities. *Waste Manag.* **35**, 329–342 (2015). [doi:10.1016/j.wasman.2014.10.006](https://doi.org/10.1016/j.wasman.2014.10.006) [Medline](#)
42. K. L. Law, Plastics in the marine environment. *Ann. Rev. Mar. Sci.* **9**, 205–229 (2017). [doi:10.1146/annurev-marine-010816-060409](https://doi.org/10.1146/annurev-marine-010816-060409) [Medline](#)
43. C. Wiedinmyer, R. J. Yokelson, B. K. Gullett, Global emissions of trace gases, particulate matter, and hazardous air pollutants from open burning of domestic waste. *Environ. Sci. Technol.* **48**, 9523–9530 (2014). [doi:10.1021/es502250z](https://doi.org/10.1021/es502250z) [Medline](#)
44. N. Reyna-Bensusan, D. C. Wilson, P. M. Davy, G. W. Fuller, G. D. Fowler, S. R. Smith, Experimental measurements of black carbon emission factors to estimate the global impact of uncontrolled burning of waste. *Atmos. Environ.* **213**, 629–639 (2019). [doi:10.1016/j.atmosenv.2019.06.047](https://doi.org/10.1016/j.atmosenv.2019.06.047)
45. C. A. Velis, Circular economy and global secondary material supply chains. *Waste Manag. Res.* **33**, 389–391 (2015). [doi:10.1177/0734242X15587641](https://doi.org/10.1177/0734242X15587641) [Medline](#)
46. R. Tramoy, J. Gasperi, R. Dris, L. Colasse, C. Fisson, S. Sananes, V. Rocher, B. Tassin, Assessment of the plastic inputs from the Seine Basin to the sea using statistical and field approaches. *Front. Mar. Sci.* **6**, 151 (2019). [doi:10.3389/fmars.2019.00151](https://doi.org/10.3389/fmars.2019.00151)
47. T. van Emmerik, M. Loozen, K. van Oeveren, F. Buschman, G. Prinsen, Riverine plastic emission from Jakarta into the ocean. *Environ. Res. Lett.* **14**, 084033 (2019). [doi:10.1088/1748-9326/ab30e8](https://doi.org/10.1088/1748-9326/ab30e8)
48. G. F. Schirrinzi, M. Köck-Schulmeyer, M. Cabrera, D. González-Fernández, G. Hanke, M. Farré, D. Barceló, Riverine anthropogenic litter load to the Mediterranean Sea near the metropolitan area of Barcelona, Spain. *Sci. Total Environ.* **714**, 136807 (2020). [doi:10.1016/j.scitotenv.2020.136807](https://doi.org/10.1016/j.scitotenv.2020.136807) [Medline](#)

## ACKNOWLEDGMENTS

We thank the anonymous peer reviewers for their suggestions which improved this manuscript. M. Hestin, L. Peake, G. Keoleian, J. Abbot, J. Muncke, and D. Mihaylova, reviewed drafts of the supplementary materials and results. We are indebted to non-authoring members of the expert panel: D. Russo, E. Moss, J. Boughton, C. Lao, D. Lerario, M. Gadgil, and M. Da, for contributing their knowledge and sharing data with this research. We are also indebted to A. Morlet, R. Opsomer, S. Defruyt, and S. Wingstrand from the Ellen MacArthur Foundation for significant contribution on knowledge and data on the plastic system and circular economy. We are grateful to J. Calcagno, J. Mandirala, J. Sirko, M. Freeman, Z. Celik, H. Watson, G. Murphy, B. Towner Levine, M. Mahoney, E. Hogan, M. Wissner, L. Mudge, A. Valdivia, and O. Tzadik, M. Ochocki and B. Frinault for conducting accuracy checks of raw data inputs, the assembly of model input files, code review, and fact checks of the supplementary material. We also thank I. Issifu and B. Frinault for reviewing manuscript drafts, C. Gonzales and A. Syed for help with data collection and curation, and P. Edwards for managing an informal review process. K. Gronsky assisted in the design of the research and provided project management support. R-M. Paquin assisted in formatting figures and references. SYSTEMIQ provided generous in-kind support to this research. **Funding:** This project was funded by The Pew Charitable Trusts. **Author contributions:** W.W.Y.L., J. E.P., Y.S., and M.R.S. conceived the study and led the overall conceptual development of this work. This research represents the equal effort and intellectual contribution among research groups at four key institutions: The Pew Charitable Trusts, SYSTEMIQ, University of Oxford, and University of Leeds, led respectively by W.W.Y.L., Y. S., R.M.B., and C.A.V. The P<sub>2</sub>O model was developed by R.M.B., and R.M.B. ran all model simulations. J.E.P. and R.M.B. processed model outputs. C.A.V. and E.R.C. provided thought leadership and expertise on solid waste management systems and practices. E.K. was a key contributor to the recycling scenarios, especially their economics and sensitivities in future changes. J.K., A.C.C., E.F. and S.B. provided the expertise in reduce and substitute interventions to select material innovations and new delivery models as well as to assess their applicability

across scenarios. J.P. prepared the figures. Y.S., J.K., E.R.C., E.J., L.K., M.E.A., M.P., T.D.P., C.G., A.S. and D.F. collected and curated the data. Working groups were led by J.K., E.R.C., M.E.A., D.F., and E.J. Manuscript drafting was led by W.W.Y.L. with significant review and editing support from J.E.P., R.M.B., M.B.M., L.G., B.D., E.R.C. and C.A.V. with further input from all authors. Technical appendices were prepared by W.W.Y.L., Y.S., R.M.B., E.R.C., J.K., C.A.V., M.B.M., E.J., K.K.H., M.P., K.L., J.E.P., coordinated by K.K.H. Data-checking of the model data inputs, including associated software development, was conducted by E.N. and M.O. and managed by M.B.M. **Competing interests:** None declared. **Data and materials availability:** The Matlab code and input files used to run the P<sub>2</sub>O model and generate the results of this study and the resulting output files are available at <https://dx.doi.org/10.5281/zenodo.3929470>.

## SUPPLEMENTARY MATERIALS

[science.sciencemag.org/cgi/content/full/science.aba9475/DC1](https://science.sciencemag.org/cgi/content/full/science.aba9475/DC1)

Materials and Methods

Figs. S1 to S6

Tables S1 to S75


References (49–219)

20 January 2020; accepted 2 July 2020

Published online 23 July 2020


[10.1126/science.aba9475](https://doi.org/10.1126/science.aba9475)






**Fig. 1. Annual rates of plastic pollution entering the environment estimated from 300 Monte Carlo simulations. (A)** Time series of plastic pollution entering aquatic and terrestrial ecosystems (Mt/y  $\pm$  95% CI) by scenario, 2016 – 2040. Scenarios: ‘Business as Usual’ (BAU), ‘Collect and Dispose’ scenario (CDS), ‘Recycling’ scenario (RES), ‘Reduce and Substitute’ scenario (RSS), and ‘System Change’ scenario (SCS). Plastic pollution rates for all scenarios between 2016 and 2020 are identical. The black point estimate in 2040 represents the annual rate of plastic pollution assuming global commitments to reduce plastic use and increase recycling announced before June 2019 are implemented prior to 2040. A time series for this scenario is not presented because timelines for implementation are unknown. **(B)** Kernel density estimates for plastic pollution (Mt) in 2040 by scenario. Boxplots of plastic pollution entering **(C)** aquatic and **(D)** terrestrial ecosystems by scenario for beginning, middle, and end years of scenario implementation.





**Fig. 2.** Fate for all municipal solid waste plastic, 2016-2040, under the 'System Change' scenario (SCS). (A) Annual mass of plastic (Mt/y) for each of five end-of-life fates. (B) Mass of plastic utility (Mt/y) addressed per modeled intervention in 2040, following 20 years of SCS implementation, organized by end of life fate. NDM = new delivery model. P2F chemical = plastic to fuel chemical conversion. P2P chemical = plastic to plastic chemical conversion. Incineration ER = Incineration with energy recovery. Aquatic poll. = plastic pollution into aquatic systems. Terrestrial poll. = plastic pollution into terrestrial systems.



**Fig. 3.** Present value costs for the management (i.e., collection, sorting, recycling, and disposal) of plastic municipal solid waste by scenario, 2016-2040. Costs (Billion 2018 USD  $\pm$  95% CI) are calculated assuming 3.5% discount rate and are net of revenues associated with the sale of recycled plastic feedstock and electricity generated from plastic incineration with energy recovery. Scenarios: Business as Usual' (BAU), 'Collect and Dispose' scenario (CDS), 'Recycling' scenario (RES), 'Reduce and Substitute' scenario (RSS), and 'System Change' scenario (SCS).





Fig. 4. Cumulative mass of plastic municipal solid waste (MSW), 2016 – 2040 (Mt ± 95% CI) polluting (A) aquatic, and (B) terrestrial systems by scenario and plastic type for years 2016-2040. (C) Cumulative mass of plastic MSW recycled for each of four plastic types modeled. (D) Cumulative mass of non-circular plastic MSW endpoints, including solutions in the mismanaged (dumpsite, open burn), effectively disposed (landfill, incineration with energy recovery, plastic to fuel (P2F) chemical conversion), and recycling (open loop recycling) categories. Uncertainty bars for P2F chemical conversion are not visible as their endpoints do not exceed the radius of the plotted point estimate. Scenarios: Business as Usual (BAU), 'Collect and Dispose' scenario (CDS), 'Recycling' scenario (RES), 'Reduce and Substitute' scenario (RSS), and 'System Change' scenario (SCS).



**Fig. 5. Fate of plastic municipal solid waste (MSW) by plastic type under the 'System Change' Scenario (SCS).** (A) Proportion of MSW ( $\pm 95\%$  CI) produced in 2040 absorbed by each of three recycling solutions and the dispose and mismanaged end-of-life categories. Even under SCS, few effective solutions are implemented to manage primary microplastics. The proportion of plastic pollution ( $\pm 95\%$  CI) entering (B) global aquatic and (C) terrestrial systems by plastic type, 2016 – 2040.

**Table 1. Plastic mass; percent of total plastic demand under different end of life fates for year 2016 and for year 2040 under the ‘Business as Usual’ (BAU) and ‘System Change’ scenarios (SCS); and percent change in plastic mass under different end of life fates for SCS in 2040 relative to 2016 and BAU in 2040.**

End of life fate	Plastic mass (Mt/y) 95% CI			Fate as % plastic demand 95% CI			SCS 2040% change 95% CI	
	2016	BAU 2040	SCS 2040	2016	BAU 2040	SCS 2040	2016	BAU 2040
Reduction	0 0, 0	0 0, 0	130 110, 150	0 0, 0	0 0, 0	31 28, 33	—	—
Substitution	0 0, 0	0 0, 0	71 62, 81	0 0, 0	0 0, 0	17 15, 18	—	—
Recycling	31 26, 32	55 46, 63	84 75, 93	14 12, 15	13 11, 15	20 18, 21	170 140, 200	54 46, 61
Disposal	97 83, 97	140 120, 150	100 89, 110	44 39, 45	32 28, 33	24 22, 26	3.5 3.3, 3.8	–26 –24, –28
Mismanaged	91 84, 100	240 220, 260	44 40, 49	42 41, 47	56 53, 59	10 9.4, 12	–51 –48, –54	–81 –76, –87
Open burn*	49 40, 60	130 110, 160	23 18, 29	54 42, 63	56 44, 65	53 41, 65	–53 –45, –61	–82 –70, –95
Dumpsite*	12 7.4, 21	25 14, 41	3.2 1.5, 5.0	13 8.2, 22	11 5.9, 17	7.3 3.3, 11	–74 –49, –99	–87 –54, –120
Aquatic pollution*	11 9.0, 14	29 23, 37	5.3 3.8, 7.0	12 9.8, 14	12 9.8, 15	12 9.0, 15	–52 –43, –60	–82 –68, –95
Terrestrial pollution*	18 13, 25	52 34, 70	12 7.8, 18	20 13, 27	22 14, 29	28 18, 39	–33 –23, –42	–76 –55, –97

\*Components of the mismanaged end of life fate. These categories sum to the total for mismanaged waste.



## Evaluating scenarios toward zero plastic pollution

Winnie W. Y. Lau, Yonathan Shiran, Richard M. Bailey, Ed Cook, Martin R. Stuchtey, Julia Koskella, Costas A. Velis, Linda Godfrey, Julien Boucher, Margaret B. Murphy, Richard C. Thompson, Emilia Jankowska, Arturo Castillo Castillo, Toby D. Pilditch, Ben Dixon, Laura Koerselman, Edward Kosior, Enzo Favoino, Jutta Gutberlet, Sarah Baulch, Meera E. Atreya, David Fischer, Kevin K. He, Milan M. Petit, U. Rashid Sumaila, Emily Neil, Mark V. Bernhofen, Keith Lawrence and James E. Palardy

published online July 23, 2020

### ARTICLE TOOLS

<http://science.sciencemag.org/content/early/2020/07/22/science.aba9475>

### SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2020/07/22/science.aba9475.DC1>

### REFERENCES

This article cites 114 articles, 6 of which you can access for free  
<http://science.sciencemag.org/content/early/2020/07/22/science.aba9475#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

---

*Science* (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2020, American Association for the Advancement of Science